



Received on 11 May 2018; received in revised form, 26 June 2018; accepted, 04 July 2018; published 01 January 2019

ANTITUBERCULAR POTENTIAL OF *DENDROPHTHOE FALCATE* (L.) AND *TRIDAX PROCUMBENS* (L.) PLANTS EXTRACTS AGAINST H37Rv STAIN OF *MYCOBACTERIA TUBERCULOSIS*

V. C. Bhagat ^{*1} and M. S. Kondawar ²

Department of Pharmaceutical Chemistry ¹, Rajgad Dnyanpeeth's College of Pharmacy, Tal- Bhor, Pune - 412206, Maharashtra, India.

Department of Quality Assurance ², Appasaheb Birnale College of Pharmacy, Sangli - 416416, Maharashtra, India.

Keywords:

Dendrophthoe falcate,
Tridax procumbens, HPTLC,
GC-MS, MABA, Antituberculosis

Correspondence to Author:

Vishwas Chandrakant Bhagat

Assistant Professor,
Department of Pharmaceutical
Chemistry, Rajgad Dnyanpeeth's
College of Pharmacy, Tal- Bhor, Pune
- 412206, Maharashtra, India.

E-mail: vishwasbhagat@rediffmail.com

ABSTRACT: The increasing incidence of MDR and XDR tuberculosis worldwide highlight the urgent need to search for newer anti-tuberculosis molecules. This research article presents a phytochemical study and antitubercular potential of methanol: water (MW), ethanol: water (EW) and dichloromethane: methanol (DM) of extracts of *Dendrophthoe falcata* (L.) and *Tridax procumbens* (L.) tested against of H37Rv *Mycobacterium tuberculosis* using microplate alamar blue assay (MABA) method. Preliminary phytochemicals studies and HPTLC fingerprint analysis revealed the presence of phytochemical like alkaloids, flavonoids, saponins, tannins, a phenolic group, glycosides, terpenoids with different R_f values. Fractions of active extracts analyzed by GC-MS shows a finding of probable phytoconstituents. Findings are useful to establishing standards for identification, purity, quality of the plant. The ethanol: water and methanol: water extracts of *Tridax procumbens* and *D. falcata* exhibited significant anti-tuberculosis activity with the MIC values of 0.8 µg/ml, 6.25 µg/ml compared to standard drug Pyrazinamide, Ciprofloxacin and Streptomycin with the MIC values of 3.125 µg/ml 6.25 µg/ml using MABA respectively against *Mycobacterium tuberculosis* (H37Rv strain) ATCC no.-27294. The presence of flavonoids, tannins, the phenolic group may contribute to the observed anti-tubercular activity. The study demonstrated that extract from *T. procumbens* and *D. falcata* could be evaluated further that might provide compounds for developing a new drug to control *M. tuberculosis*.

INTRODUCTION: Infectious diseases represent a critical issue for health and are the major cause of morbidity and mortality worldwide. The impact is even greater in developing countries due to unavailability of medicine.

Tuberculosis remains a global public health problem in developing countries. Due to the global emergence of multidrug-resistant (MDR) and extensively drug resistant (XDR) strains of *M. tuberculosis* ¹.

There is an urgent need to develop new drugs and strategies to fight TB. The World Health Organisation (WHO) estimated that approximately 80% of the world population relies mainly on traditional medicines, mainly plant drugs in their health care. The last few decades have witnessed a substantial increase in the investigation of

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.10(1).251-59
	The article can be accessed online on www.ijpsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.10(1).251-59	

medicinal plants for their biological efficacy in the treatment of various disorders. Over the years, some improved and high throughput techniques towards screening of anti-mycobacterial agents have been developed^{2,3}. Several methods exist for testing the antitubercular potential of plant extracts such as fluorescence-based testing on the Bactec MGIT960 system, use of redox indicator dyes such as alamar blue or resazurin and MTT, using colony forming units (CFU) on solid agar plates. Techniques such as the agar diffusion and broth dilution method have been used^{4,5}. The microplate alamar blue assay (MABA) is a colorimetric oxidation-reduction based assay. It is a non-radiometric, rapid, high-throughput and comparatively low-cost assay producing results with a high degree of confidence^{6,7,8}.

Dendrophthoe falcata (L.f) Ettingsh is highly specialized perennial flowering plant adapted to parasitic life on aerial parts of their hosts. The whole parasitic plant is used in indigenous system of medicine as cooling, bitter tonic, astringent, aphrodisiac, narcotic, diuretic, pulmonary tuberculosis, asthma, menstrual disorders, swelling wounds, ulcers, renal and vesicle calculi and violated conditions of kapha and pitta⁹. *D. falcata* belonging to the family Loranthaceae is an angiosperm hemiparasitic plant was most frequently observed on many host plants comprises of 20 species, and about 7 species are found in India. The medicinal properties of *D. falcata* are generally influenced by the host plant for example when grown on *Calotropus gigantea* (Linn.) (Asclepiadaceae) the parasitic plant is considered useful for improving cognitive function, on *Tamarindus indicus* L. (Fabaceae) it is used to treat impotence and on *S. robusta* (Dipterocarpaceae) it is used to treat paralysis¹⁰. *D. falcata* bark juice/decoction used in menstrual troubles and asthma while its paste is applied on boils, setting dislocated bones and extracting pus.

The decoction of the whole plant is used to treat joint pains¹¹. Quercetin, kaempferol, quercetin are the different flavonoids isolated and reported from the whole plant of *D. falcata*^{12,13}. The reported biological activities of *D. falcata* are antioxidant, anti-inflammatory¹⁴, antitumor^{15,16}, anthelmintic potential¹⁷, anti-nociceptive and general toxic studies¹⁸, anticonvulsant and muscle relaxant¹⁹,

immunomodulatory²⁰, wound healing and antimicrobial²¹, antifertility²², anticancer studies²³. *D. falcata* (Loranthaceae) a parasite on *Mangifera indica* (Anacardiaceae) potential antidiabetic²⁴, Anti-diarrhoeal and insecticidal activities²⁵.

Tridax procumbens Linn. family Compositae commonly known as 'coat buttons' because of an appearance of flowers has been extensively used Ayurveda in liver disorders²⁶. *Tridax* possesses significant anti-inflammatory, hepatoprotective, wound healing, antidiabetic activity and antimicrobial activity against both gram-positive and gram-negative bacteria^{27,28,29,30}. Flavonoids isolated and reported from whole plant^{31,32}. The leaf juice possesses antiseptic, insecticidal and parasiticidal properties and is also used to check hemorrhage from cuts, bruises and wound. Interestingly it also has a hypotensive effect and potent immunomodulating property^{33,34} — *in-vitro* radical scavaging and *in-vivo* anti-inflammatory potential³⁵. The primary reasons to select plants is their known antimicrobial properties, and other pharmacological uses as an antioxidant, immunomodulatory also used for respiratory disorders in reported ethnobotanical surveys was also documented are base for research. The study was undertaken to evaluate a phytochemical analysis and preliminary report on the antitubercular activity of *D. falcata* and *Tridax procumbens* plant extracts against M. tuberculosis (H37R_v strain) using bio-assay guided fractionation of the extracts of the leaves and fractions thereof.

MATERIALS AND METHODS:

Chemicals: Chemicals used in this study were of analytical grade, and highest purity procured from standard commercial sources Research Lab. and S. D. Fine Lab Mumbai.

Plant Collection and Identification: The *Tridax procumbens* Linn. (Asteraceae) plant and *Dendrophthoe falcata* (L.f) Etting. (Loranthaceae) samples were collected at the flowering stage from the local region, during September - November was collected from Kapurhol Kasurdi on Satara-Pune NH-4 near Pune Maharashtra (India). Both specimens plant were identified and authenticated by Botanical Survey of India (BSI) Pune, Maharashtra India having voucher specimen number VIBTRP2 & VIBDEF3 dated-15/02/2013.

Preparation of Extracts: Extracts were prepared in a sequential manner using ethanol: water, methanol: water, dichloromethane: methanol (order of increasing polarity) as solvents from of shade-dried and coarsely powdered plant material using the maceration and Soxhlet apparatus. Coarsely powdered was defatted with petroleum ether and then exhaustively extracted with a different solvent.

The methanol: water and ethanol: water extract was prepared by maceration by soaking 10g of powdered plant materials in 100 ml of solvent at room temperature for 48 h. The extract was filtered after 48 h, through a sterilized Whatman no. 1 filter paper. The extract concentrated using a rotary vacuum evaporator with the water bath set at 40 °C. Overnight UV-irradiation sterilized the dried extract. The sterile extract was transferred into a sterile lyophilization flask and frozen in a deep freezer. The extract was stored at -20 °C till bio-evaluation.

Phytochemical Analysis: The phytochemical investigation of the different extracts of *Tridax procumbens* and *Dendrophthoe falcata* was carried out with standard protocol^{36, 37}. Extracts were evaluated for physical constants³⁸.

HPTLC Study: HPTLC is an important analytical tool in the separation, identification, and estimation of various classes of natural phytoconstituents. HPTLC studies were carried out by the method of Harborne³⁹ and Wagner *et al.*,^{40, 41}. The extracts were dissolved in the respective solvents (5mg/ml). The 10µl of sample extract was applied with the help of linomat syringe using the linomat applicator 5. Solvent system *n*-hexane: toluene: ethyl acetate (2:4:1). HPTLC silica gel F254 (Merck). The plates were developed in a CAMAG chamber. CAMAG HPTLC densitometer (Scanner) used to measure absorbance mode at 254, 366 nm and 560 nm. Data integration through the software WINCATS Planar Chromatography Manager. The fingerprint so developed and R_f value were noted. Spots were visible without derivatization at 254 nm, 366 nm, and 560 nm wavelengths. A solvent system optimized for TLC study was chosen for HPTLC study.

Anti-tuberculosis Activity:

Microbial Strain for Anti-Mycobacterium tuberculosis Assay: Reference strains H37Rv

(ATCC No-27294) of *Mycobacteria tuberculosis* (Vaccine strain) used.

Microplate Alamar Blue Assay: The *M. tuberculosis* (MTB) were cultured in 7H9 medium in the presence of the plant extracts in a 96 well plate was tested at concentration 0.8, 1.6, 3.12, 6.25, 12.5, 25, 50 and 100 µg/ml. The anti-mycobacterial activity of compounds was assessed against *M. tuberculosis* using MABA. This methodology is non-toxic, uses a thermally stable reagent and shows good correlation with proportional and BACTEC radiometric method. 200µl of sterile deionized water was added to all outer perimeter wells of sterile 96 wells plate to minimized evaporation of medium in the test wells during incubation. The 96 wells plate received 100µl of the Middlebrook 7H9 broth, and serial dilution of compounds was made directly on the plate. The final drug concentrations tested were 0.2 to 100µg/ml. Plates were covered and sealed with parafilm and incubated at 37 °C for five days. After this 25µl of freshly prepared 1:1 mixture of Almar Blue reagent and 10% tween 80 was added to the plate and incubated for 24 h. A blue color in the well was interpreted as no bacterial growth, and pink color was scored as growth. Interpretations were based on the percent reduction of the dye which is directly proportional to the bacterial growth^{42, 43}. The extracts were considered active if the percent reduction value of alamar blue dye was less than that observed for the standard. Triplicate wells were maintained for each variable in every assay, and all the assays were performed thrice.

GC-MS Analysis: Fractions of active extracts analyzed by GC-MS Quadrupole Analyzer at Poona College of Pharmacy, Pune (Maharashtra) India. Shows finding some probable Phyto-constituents in the mass range 300-600 am.

RESULTS:

Physical Constants: The proximate analysis showed a satisfactory result concerning the foreign matter, moisture content, ash value, and extractive values⁴⁴. The physical constants are shown in **Table 1**.

Photochemical Screening: The extracts were screened for phytochemical constituents for the presence of saponins, tannins, alkaloids, flavonoids, a phenolic group, glycosides and reducing sugars³⁷.

The presence of alkaloids, glycosides, flavonoids, tannins, phenolic, xanthenes, quinones, sterols, triterpenoids, etc is mostly responsible for the anti-tubercular activity proven in Ayurveda ⁴⁵. The results are shown in **Table 2**.

HPTLC Fingerprint Studies: The study revealed that *Tridax procumbens* (Linn.) and *Dendrophthoe falcata* (Linn.) showed best results in *n*-hexane: toluene: ethyl acetate (2: 4: 1) solvent system for TPEW, DFMW, and DFDM extracts.

TABLE 1: EVALUATION OF PHYSICAL CONSTANTS OF POWDERED *TRIDAX PROCUMBENS* (L.) AND *DENDROPHTHOE FALCATA* (L.)

S. no.	Evaluation parameter	Value (%)	
		<i>Tridax procumbens</i> (L.)	<i>Dendrophthoe falcate</i> (L.)
1	Foreign matter	1	1
2	Moisture content	10.6	9.5
3	Total ash value	12.7	11.9
4	Water soluble ash value	3.5	3.2
5	Acid-insoluble ash value	8.0	9.1
6	Water soluble extractive value	4.89	3.87
7	Chloroform soluble extractive value	0.4	0.5
8	Methanol soluble extractive value	6.66	6.73
9	Ethanol soluble extractive value	4	5.2

TABLE 2: PRELIMINARY PHOTOCHEMICAL SCREENING OF EXTRACTS OF *T. PROCUMBENS* (L.) AND *D. FALCATA* (L.)

Phyto-constituents	<i>Tridax procumbens</i> extracts			<i>Dendrophthoe falcata</i> extracts		
	Methanol: Water	Ethanol: Water	Dichloromethane: Methanol	Methanol: Water	Ethanol: Water	Dichloromethane: Methanol
Alkaloids	-	+++	++	+	-	+
Flavonoids	++	+++	+++	++	+++	+++
Phenolic groups	+	+	-	-	-	-
Saponin Glycosides	++	+++	+	++	+	-
Tannins	+++	+++	+++	+++	+++	+++
Steroids	++	+++	++	++	++	++
Carbohydrates	+	+	+	+	+	+
Terpenoids	+	++	+	+	++	+
Cardiac Glycosides	-	-	-	+	+	+
Reducing sugar	+	-	++	+	-	+
Anthroquinone Glycosides	-	-	-	-	-	+

+ indicates presence - indicates absence.




FIG. 1: CHROMATOGRAM OF ETHANOL: WATER EXTRACT OF *TRIDAX PROCUMBENS* MEASURED AT 366nm




FIG. 2: CHROMATOGRAM OF METHANOL: WATER EXTRACT OF *D. FALCATA* (L.) MEASURED AT 366 nm




FIG. 3: CHROMATOGRAM OF DICHLOROMETHANE: METHANOL EXTRACT OF *D. FALCATA* (L.) MEASURED AT 560 nm

After scanning and visualizing the plates in absorbance mode at both 366 nm, 560 nm. The HPTLC images shown that all sample constituents were separated without any tailing and diffuseness. As a shown in **Fig. 1**, **Fig. 2** and **Fig. 3**.

Anti-Tuberculosis Evaluation: The anti-microbial effects of the extracts evaluated by using microplate alamar blue assay (MABA) against *M. tuberculosis* (H37Rv strain) ATCC No-27294, were evaluated at the Department of Microbiology,

Maratha Mandal's NGH Institute of Dental Sciences and Research Centre, Belgaum - 590010, India.

The method applied is similar to that reported by Maria and Lourenco⁴³. Pyrazinamide (MIC- 3.125 µg/ml), Ciprofloxacin (MIC-3.125 µg/ml) and Streptomycin (MIC-6.125 µg/ml) were used as reference standard to evaluate the potency of the extracts. As shown in **Table 3** and **Fig. 4**, **Fig. 5** and **Fig. 6**.

TABLE 3: SHOWING COMPARATIVE ANTI-TUBERCULOSIS SCREENING RESULTS BY MIC METHOD

Code no.	Compounds / Extracts	MIC values µg/ml	Code no.	Compounds / Extracts	MIC values µg/ml
A1	DFDM	12.5	Std 1	Pyrazinamide	3.125
A2	DFEW	25	Std 2	Ciprofloxacin	3.125
A3	TPMW	25	Std 3	Streptomycin	6.25
B1	TPDM	25			
B2	TPEW	0.8			
B3	DFMW	6.25			

T.P -*Tridax procumbens* and D.F- *Dendrophthoe falcata*




FIG. 4: MIC µg/ml OF TESTED EXTRACTS AGAINST H37Rv OF MYCOBACTERIA TUBERCULOSIS

*DFDM- *Dendrophthoe falcata* dichloromethane-methanol, DFEW - *Dendrophthoe falcata* ethanol-water, TPMW- *Tridax procumbens* methanol-water, TPDM- *Tridax procumbens* dichloromethane-methanol, TPEW- *Tridax procumbens* ethanol-water, DFMW- *Dendrophthoe falcata* methanol-water




FIG. 5: MICROPLATE ALAMAR BLUE ASSAY 96 WELL PLATE FOR ACTIVITY OF PLANT EXTRACTS




FIG. 6: MICROPLATE ALAMAR BLUE ASSAY 96 WELL PLATE FOR ACTIVITY OF STD

GC-MS Analysis: GC-MS analysis of *Tridax procumbens* (L.) active ethanol-water extract shows

finding number of probable phytoconstituents in Mass range 300-600 amu **Fig. 7 - 12**.




FIG. 7: GC-MS SPECTRA OF PHYTOCONSTITUENT-1




FIG. 8: GC-MS SPECTRA OF PHYTOCONSTITUENT-2




FIG. 9: GC-MS SPECTRA OF PHYTOCONSTITUENT-3




FIG. 10: GC-MS SPECTRA OF PHYTOCONSTITUENT-4




FIG. 11: GC-MS SPECTRA OF PHYTOCONSTITUENT-5




FIG. 12: GC-MS SPECTRA OF PHYTOCONSTITUENT-6

DISCUSSION: The Antimycobacterial activities of extracts were evaluated by using microplate alamar blue assay (MABA). In the present study, the TPEW extract exhibited significant anti-tuberculosis activity with the MIC values of 0.8 μ g/ml against H37Rv of *M. tuberculosis*. The DFMW extract exhibited significant anti-tuberculosis activity with the MIC values of 6.25 μ g/ml against H37Rv of *M. tuberculosis*.

DFDM extract shows moderate anti-tuberculosis activity with the MIC values of 12.50 μ g/ml. The DFEW extract and TPMW, TPDM extract displayed a weak activity against H37RV of *M. tuberculosis* strain compare to standard drugs as Pyrazinamide, Ciprofloxacin, and Streptomycin with the MIC values of 3.125 μ g/ml, 3.125 μ g/ml and 6.25 μ g/ml using microplate alamar blue assay

(MABA). As many different methods are available to evaluate antituberculosis activity, no specific cut-off value has been established for reference to analyze the anti-tuberculosis activity⁴⁶. Values show that out of the six, two extracts was found to be more potent than std - drugs as Pyrazinamide, Ciprofloxacin, and Streptomycin.

The preliminary phytochemical screening of various extracts shows the presence of alkaloids, flavonoids, saponins, tannins, a phenolic group, glycosides, terpenoids, reducing sugars. On investigation of extracts having significant anti-tuberculosis activity is revealed the presence of alkaloids, flavonoids, saponins, tannins, a phenolic group, glycosides, terpenoids. Flavonoids may act by depolarization of membrane and inhibition of DNA, RNA and proteins synthesis.

It may be reduced the bacterial cell density rapidly and caused lysis⁴⁷. The efficacy of extracts could be due to the interplay of active constituents present, leading to better activity. It has been demonstrated that different constituents of crude extracts act through different mechanisms. Tannin is used as antimicrobial growth-promoting factor (AGP) in the sub-therapeutic dose for long periods is particularly favorable for the selection of antimicrobial resistant microorganism. Chalcones as a flavonoids, 1-(2-hydroxyphenyl)-3-(3-chlorophenyl)-2-propane-1-one and 1-(2-hydroxyphenyl)-3-(4-iodophenyl)-2-propane-1-one, showed inhibition (90%) of *Mycobacterium tuberculosis* H37Rv. Some chalcone like a compound with heterocyclic ring showed even higher inhibition (95%) as anti-tubercular agent⁴⁸.

The HPTLC fingerprint analysis chromatograms developed are particular with the finalized solvent system, *n*-hexane: toluene: ethyl acetate in the ratio of (2: 4: 1). HPTLC profiling of the extract confirms about the presence of various phytochemicals. R_f value % area can serve as an improved tool for the extract/subfraction standardization.

The present study gives enough information regarding various phytoconstituents present in the methanol: water, ethanol: water, dichloromethane: methanol extract of *Tridax procumbens* (Linn.) and *Dendrophthoe falcata* (Linn.) also helps in generating the basis for the quality control, correct identification, and standardization.

The HPTLC finger print of ethanol: water extract of *Tridax procumbens* (L.) clearly showed signified existence of 12 phytoconstituents whose R_f values ranged from 0.06 to 0.87 eluting out at 366 nm With R_f values 0.87, 0.82, 0.76, 0.70 were found to be leading as the percentage area was more, *i.e.* 25.74%, 5.32%, 14.18%, and 6.34% respectively. Fractions of active extracts analyzed by GC-MS shows finding a number of probable phytoconstituents. The study revealed that *T. procumbens* and *D. falcata* was rich in secondary metabolites particularly tannins and flavonoids which are may be responsible for the antitubercular activity. It identifies *T. procumbens* and of *Dendrophthoe falcata* have a promising antitubercular activity.

CONCLUSION: The study indicated that selected *Tridax procumbens* (L.) and of *Dendrophthoe falcata* (L.) leaf extracts exhibited potential anti-tubercular activity. The present investigation may be the focus of further phytochemical research of *Tridax procumbens* and *Dendrophthoe falcata* to identify and isolate the constituents responsible for antitubercular activity are being undertaken, along with the exploration of mechanisms of action and contribute greatly to the development new phytomolecule against tuberculosis.

ACKNOWLEDGEMENT: The authors are thankful to the head of the Institutions for providing the necessary facilities, thankful to Botanical Survey of India (BSI) Pune, India. We thanks to Dr. Kishor Bhat, Department of Microbiology, Maratha Mandal's NGH Institute of Dental Sciences and Research Centre, Belgaum-590010, India. For anti-tubercular activity, the authors are thankful to Poona College of Pharmacy, Pune (Maharashtra) India. For GC-MS instrumentation facility.

CONFLICT OF INTEREST: The authors declare no conflict of interest.

REFERENCES:

- 1 Singh C, Singh SK, Nath G and Rai NP: Antimycobacterial activity of *Piper longum* L. fruit extracts against multi-drug resistant *Mycobacterium spp.* International Journal of Phytomedicine 2011; 3: 353-361.
- 2 Kumar V, Patel S and Jain R: New structural classes of antituberculosis agents. Med Res Rev 2018; 38(2): 684-740.
- 3 Forbes L, Ebsworth-Mojica K, Di-Dono L, Li SG, Freundlich JS and Connell N: A high throughput screening assay for anti-mycobacterial small molecules based on adenylate kinase release as a reporter of cell lysis. PLoS One 2015; 10(6): 1371.
- 4 Gautam R, Saklani A and Jachak SM: Indian medicinal plants as a source of antimycobacterial agents. Journal of Ethnopharmacology 2007; 110: 200-234.
- 5 Kumar KJ, Devi Prasad AG and Chaturvedi V: Phytochemical screening of five medicinal legumes and their evaluation for *in-vitro* anti-tubercular activity. AYU 2014; 35(1): 98-101.
- 6 Collins LA and Franzblau SG: Microplate alamar blue assay vs. BACTEC 460 for high-throughput screening of compounds against *Mycobacterium tuberculosis* and *Mycobacterium avium*. Antimicrob Agents Chemother 1997; 4: 1004-1009.
- 7 Kohli A, Bashir G, Fatima A, Jan A, Wani N and Ahmad J: Rapid drug-susceptibility testing of *Mycobacterium tuberculosis* clinical isolates to first-line antitubercular drugs by nitrate reductase assay a comparison with proportion method. Inter Journal of Mycobacteriology 2016; 5: 469-474.

- 8 Wahyuningrum R, Ritmaleni, Irianti T, Wahyuono S, Kaneko T and Nuryastuti T: Antituberculosis activity of Brotowali (*Tinospora crispa*) extract and fractions against *Mycobacterium tuberculosis* using microplate alamar blue assay method. *Traditional Medicine Journal* 2017; 22(2): 124-130.
- 9 Nadkarni KM: *Indian material medica* 1993; 1: 1276-1277.
- 10 Pattanayak SP, Mazumder PM and Sunitha P: *D. falcata* (L.f) Ettingsh: A consensus review. *Pharmacog* 2008; 2: 359-368.
- 11 Jagtap SD, Deokule SS and Bhosle SV: Some unique ethnomedicinal uses of plants used by the Kokru tribe of Amravati district of Maharashtra, India. *Journal of Ethnopharmacol* 2006; 107: 463-469.
- 12 Nair AGR and Krishnakumary P: Flavonoids from *Dendrophthoe falcata* Ettingsh growing on different host plants. *Indian J. Chem* 1998; 29: 584-585.
- 13 Mallavadhani UV: Three new pentacyclic triterpenes and some flavonoids from the fruits of an Indian Ayurvedic plant *Dendrophthoe falcata* and their estrogen receptor binding activity. *Chem Pharm Bull* 2006; 54: 740-44.
- 14 Patil S, Anarthe S, Jadhav R and Surana S: Evaluation of anti-inflammatory activity and *in-vitro* antioxidant activity of Indian mistletoe, the Hemiparasite *Dendrophthoe falcata* L.F. (Loranthaceae). *Iran J Pharm Res* 2011; 10(2): 253-259.
- 15 Dashora N, Sodde V, Prabhu KS and Lobo R: *In-vitro* cytotoxic activity of *dendrophthoe falcata* on human breast adenocarcinoma cells-mcf7. *International Journal of Cancer Research* 2011; 7(1): 47-54
- 16 Dashora N and Sodde V: Antitumor activity of *Dendrophthoe falcata* against Ehrlich Ascites carcinoma in Swiss albino mice. *Pharmaceutical Crops* 2011; 2: 1-7.
- 17 Raut DN, Pal SC and Mandal SC: Anthelmintic potential of *Dendrophthoe falcata* Etting. (L.F) Leaf. *International Journal of Pharmaceutical Research and Development* 2009; 6: 12-17.
- 18 Hasan M: Antioxidant, antinociceptive activity and general toxic studies of *D. falcata* and isolation of quercitrin as the major component. *Oriental Pharmacy and Wxper Mental Medicine* 2006; 6(4): 355-360.
- 19 Sinoriya P, Irchhaiya R and Kumar S: Anticonvulsant and muscle relaxant activity of the ethanolic extract of stems of *Dendrophthoe falcata* (Linn. F.) in mice. *Indian Journal of Pharmacol* 2011; 43(6): 710- 713.
- 20 Pattanayak SP and Mazumder PM: Immunomodulatory activities of *Dendrophthoe falcata* (L. f) Ettingsh in experimental animals: *in-vitro* and *in-vivo* investigations. *Journal of Scientific Research* 2011; 3(3): 619-630.
- 21 Pattanayak SP: Wound healing, antimicrobial and antioxidant potential of *Dendrophthoe falcata* (L.f) Ettingsh. *Journal of Ethnopharmacology* 2008; 120(2): 241-247.
- 22 Pattanayak SP and Mazumder PM: Effect of *D. falcata* (L.F) Ettingsh on the female reproductive system in Wistar rats: a focus on anti-fertility efficacy. *Epub* 2009; 80(3): 314-320.
- 23 Kodithala S, Yoganandam GP and Kiranmai M: Pharmacognostical phytochemical and anticancer studies of *Dendrophthoe falcata* (L.f.) ettingsh. (Loranthaceae) growing on the host plant *Azadirachta indica* (Meliaceae). *Int J Pharm Bio Sci* 2013; 4(2): 1010-1018.
- 24 Anarthe SJ, Bhalke RD, Jadhav RB and Surana SJ: Anti-diabetic activity of stems of hemiparasite *Dendrophthoe falcata* Linn. parasitic growing on *Mangifera indica* Dhaka Univ. *J. Pharm. Sci* 2008; 7(2): 177-180.
- 25 Haque A, Zaman A, Hossain M, Sarker I and Islam S: Evaluation of anti-diarrhoeal and insecticidal activities of ethanol extract and its fractions of *Dendrophthoe falcata* (L.) leave. *IJPSR* 5(9): 3653-3663.
- 26 Saraf S and Dixit VK: Hepatoprotective activity of *Tridax procumbens* Part -II, *Fitoterapia*, 1991; 62: 534-536.
- 27 Ravikumar V, Shivashangari KS and Devaki T: Hepatoprotective activity of *Tridax procumbens* against dgalactosamine / lipopolysaccharide-induced hepatitis in rats. *Journal of Ethnopharmacology* 2005; 101: 55-60.
- 28 Raina R, Prawez S, Verma PK and Pankaj NK: Medicinal plants and their role in wound healing. *Vet Scan* 2008; 3(1): 221-224.
- 29 Bhagwat DA, Killedar SG and Adnaik RS: Antidiabetic activity of leaf extract of *Tridax procumbens*. *Int. J. Green Pharma* 2008; 2: 126- 128.
- 30 Mahato RB and Chaudhary RP: Ethnomedicinal study and antibacterial activities of selected plants of Palpa District, Nepal. *Scientific World* 2005; 3(3): 26-31.
- 31 Saxena VK and Albert S: b-Sitosterol-3-O-b-Dxylopyranoside from the flowers of *Tridax procumbens* Linn. *J. Chem. Sci* 2005; 117(3): 263-266.
- 32 Ali M, Ravinder E and Ramchandram R: Phytochemical communication a new flavonoid from the aerial parts of *Tridax procumbens*. *Fitoterapia* 2001; 72: 313-315.
- 33 Tiwari U, Rastogi B, Singh P, Saraf DK and Vyas SP: Immunomodulatory effects of aqueous extract of *Tridax procumbens* in experimental animals. *Journal of Ethnopharmacology* 2004; 92: 113-119.
- 34 Oladunmoye MK: Immunomodulatory effects of ethanolic extract of *Tridax procumbens* on Swiss albino rat's orogastrically dosed with *Pseudomonas aeruginosa* (NCIB 950). *International Journal of Tropical Medicine* 2006; 1(4): 152-155.
- 35 Nia R, Paper DH, Essien EE, Oladimeji OH, Iyadi K and Franz G: Investigation into *in-vitro* radical scavaging and *in-vivo* anti-inflammatory potential of *Tridax procumbens*. *Nigerian Journal of Physiological Science* 2003; 18(1-2): 39-43.
- 36 Quality Control Method for Medicinal Plant Materials; WHO, Geneva 1998: 1-15.
- 37 Sharma MK and Sharma S: Phytochemical and Pharmacological screening of combined *Mimosa pudica* Linn. and *Tridax procumbens* for *in-vitro* antimicrobial activity. *Int. J. Microbiol. Res* 2010; 1 (3): 171-174.
- 38 Khandelwal KR: Techniques and experiments, practical pharmacognosy. Nirali Prakashan, Pune, Edition 17th, 2007: 149-156.
- 39 Harborne JB: *Phytochemical methods*. Chapman and Hall, London, Edition 3rd, 1998.
- 40 Wagner H and Baldt S: *Plant drug analysis*. Springer, Edition 2nd, 1996.
- 41 Chandrakasan L and Neelamegam R: HPTLC analysis of coumarin profile in the leaf and bark samples of *Loranthus longiflorus* Desr. (Synonym – *Dendrophthoe falcata* (L.f.) Ettingsh) collected from two host trees. *Journal of Medicinal Plants Studies* 2017; 5(1): 135-139.
- 42 Collins LA and Franzblau SG: Microplate alamar blue assay vs BACTEC 460 for high-throughput screening of compounds again *Mycobacterium tuberculosis* and *Mycobacterium avium*. *Antimicrob. Agents Chemother* 1997; 41: 1004-1009.
- 43 Lourenco MCS, Desouza MVN and Pinheiro AC: Evaluation of the anti-Tubercular activity of nicotinic and isoniazid analogues. *ARKIVOC* 2007; 181-191.
- 44 Ejoba R: Phytochemical constituents of some leaves extract of *Aloe vera* and *Azadirachta indica* plant species.

- Global Advanced Research Journal of Environmental Science and Toxicology 2012; 1(2):14-17.
- 45 Arya V: A review on anti-tubercular plants. International Journal of Pharm Tech Research 2011; 3(2): 872-880.
- 46 Luo X and Pires D: Antimycobacterial evaluation and preliminary phytochemical investigation of selected medicinal plants traditionally used in Mozambique. Journal of Ethnopharmacology 2011; 137: 114-120.
- 47 Dzoyem JP, Hamamoto H and Ngameni B: Antimicrobial action mechanism of flavonoids from *Dorstenia species*. Drug Discoveries and Therapeutics 2013; 7(2): 66-72.
- 48 Lin YM and Zhou Y: Chalcones and flavonoids as an anti-tuberculosis agent. Bioorg Med Chem 2002; 10: 2795-02.

How to cite this article:

Bhagat VC and Kondawar MS: Antitubercular potential of *Dendrophthoe falcate* (L.) and *Tridax procumbens* (L.) plants extracts against H37Rv stain of *Mycobacteria tuberculosis*. Int J Pharm Sci & Res 2019; 10(1): 251-59. doi: 10.13040/IJPSR.0975-8232.10(1).251-59.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Play store)